



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

wonderful marine aviary—second to none of the natural features of California.” In 1884 it is said 300,000 eggs were gathered and the market was glutted, while the present year only 91,740 have been taken. Comparatively few birds are allowed to breed and such merciless persecution can but result in extinction. The commercial value of these rookeries will doubtless prevent their protection from purely sentimental grounds, but if it can be shown that the present course will end in the destruction of the egg industry, it might be possible to secure the enactment of a law which would protect the birds for at least the latter half of the nesting season.—F. M. C.

Ridgway on New Species and Subspecies of Birds¹.—In the three papers here cited, none of which bears date of publication, Mr. Ridgway describes *Geothlypis flaveolatus* from near Tampico on the Gulf coast of Mexico, a form which, strangely enough, is most closely related to *G. beldingi* of Lower California; *Geospiza pachyrynchos*, *G. fatigata*, *Camarhynchus bindloei*, *C. compressirostris*, and *C. incertus* from the apparently exhaustless Galapagos, and *Peucedramus olivaceus aurantiacus* from Guatemala.—F. M. C.

Oberholser on Two New Subspecies of Dryobates².—Comparison of a series of 200 Downy Woodpeckers has impressed Mr. Oberholser with the differences in size and color existing between specimens from the South Atlantic and Gulf States and those from Alaska and northern British America, and he therefore separates these extremes under the names *Dryobates pubescens meridionalis* (Swainson) and *Dryobates pubescens nelsoni* (Oberholser) respectively, leaving *Dryobates pubescens* as a transition form occupying the intervening region.—F. M. C.

Richmond on Mexican Birds³.—This is a nominal list of 58 species, with the number of specimens of each, received by the National Museum

¹ 1. Description of a New Species of Ground Warbler from Eastern Mexico. By Robert Ridgway, Curator of the Department of Birds. Proc. U. S. Nat. Mus., XVIII, p. 119.

2. Preliminary Description of some New Birds from the Galapagos Archipelago. *Ibid.*, p. 293.

3. Description of a New Subspecies of the Genus *Peucedramus*, Coues. *Ibid.*, p. 441.

² Description of Two New Subspecies of the Downy Woodpecker, *Dryobates pubescens* (Linnæus). By Harry C. Oberholser. Proc. U. S. Nat. Mus., XVIII, 1895, p. 547.

³ Partial list of Birds collected at Alta Mira, Mexico, by Mr. Frank B. Armstrong. By Charles W. Richmond, Assistant Curator of the Department of Birds. Proc. U. S. Nat. Mus., XVIII, 1896, pp. 627–632.